
Marija Radomirović¹

THE ROLE OF THE CIRCADIAN SYSTEM IN CARDIOVASCULAR DISEASES AND CLINICAL IMPLICATIONS

Abstract: Circadian rhythm, an internal 24-hour cycle biological clock, regulates important biologic functions such as metabolism, body temperature, heart and respiratory rate, circulating levels of hormones, sleep-wake pattern, behaviour. It is also a very important in regulation of the cardiovascular (CV) system components such as arterial blood pressure, pulse, endothelial function. In this comprehensive review of the literature, we will discuss how interaction between environment/behavior stressors and circadian phases modulate individual cardiovascular risk. Potential therapeutical targets will be also discussed.

Keywords: circadian rhythm; cardiovascular system; cardiovascular diseases

Introduction

It is well known that all multicellular organisms, including humans, have an internal, biological clock that regulates important biologic functions such as metabolism, body temperature, heart and respiratory rate, circulating levels of hormones, sleep/wake pattern, behavior. The biological clock provides that the body's processes are optimized at various points during a 24-hour period (1). The 2017 Nobel Prize in Physiology or Medicine is awarded to the scientists for their discoveries of genes that regulate circadian rhythm, described in 1984, in the fruit fly model (*Drosophila*). Jeffrey C. Hall and Michael Rosbash showed that clock gene, called *period*, encoding protein called PER, accumulates in the cell during the night, and degrades during the day. PER protein levels oscillate during 24h, synchronously to the circadian rhythm. A decade later, in 1994, Michael W. Young discovered *timeless*, an additional gene influencing the circadian clock, that encodes TIM protein. Upon entering the nucleus, the mentioned proteins block the activity of the period gene by a negative feedback mechanism. Young also identified the *doubletime* gene, which encodes a DBT protein

¹ Marija Radomirović, Cardiology Clinic of the University Clinical Center of Serbia, e-mail: marija3793@gmail.com.

whose role is to delay the accumulation of PER protein (2). Considering the complex and precise molecular mechanisms, the clinical significance of these findings is that chronic disturbance of the biological rhythm is associated with an increased risk for various diseases, primarily cardiovascular, then endocrinological and malignant.

Circadian rhythm

Franz Halberg, one of the founders of chronobiology, introduced the term circadian rhythm to science in 1959 (latin. *Circa* – around, and *dies* – day), which he defined as a cyclic change of a certain biological function with duration of one cycle about 24 hours, independent of external factors oscillations (Halberg, 1959). Molecular analyzes have shown that every cell in the body undergoes circadian clock with a master biological clock, that is, the circadian pacemaker, the suprachiasmatic nucleus (SCN) in the hypothalamus, which synchronizes and coordinates other oscillators in the body. Parvocellular neurons of the SCN receive information from the retina, via the retinohypothalamic pathway. In addition to numerous roles, melatonin synthesis and secretion control is a major output function of the SCN. The light falling on the retina sends signals to the pineal gland (latin. *glandula pinealis*), inhibits melatonin synthesis and stimulates wakefulness (3). Melatonin the main product of the pineal gland, and the primary hormone of the circadian system, exhibits numerous antioxidant, anti-inflammatory, antidepressant, immunomodulatory and cardioprotective effects. Biosynthesis of melatonin mainly occurs in the pineal gland. Although it is also synthesized in the retina, digestive tract, bone marrow and lymphocytes, synthesis in these tissues is not regulated by the light/dark signal. The start of melatonin secretion is usually around 09:00-10:00 p.m. with a maximum between midnight and 03:00 a.m. while the lowest concentration is in the early morning hours, between 07:00 and 09:00 a.m. (4).

The role of the circadian rhythm in the cardiovascular system

Central oscillators are present in the central nervous system, while peripheral oscillators are found in various organs, including the heart. Genes, receptors, cells, organs and physiological processes in the cardiovascular system have circadian rhythmicity. In the body, there is a constant interaction between behavioral stressors (for example, a sudden change in body position after getting up, sudden rise in blood pressure, change of night and day, food intake), circadian phases and individual cardiovascular risk. Parameters of the cardiovascular system such as arterial blood pressure, heart rate, vascular endothelium function, coagulation factors, level of circulating catecholamines and others, show physiological 24-hour variations (5).

Circadian variations in blood pressure

Blood pressure is characterized by circadian variations, with the most pronounced drop in values during the night by 10 to 20%, both in healthy individuals and in hypertensive patients. The “non-dipping” phenomenon is a drop in blood pressure during the night by less than 10% and is associated with an increased cardiovascular risk (6). Similarly, in elderly patients with hypertension, an increase in systolic blood pressure more than 10 mmHg two hours after waking up was associated with a 22% higher risk of stroke, regardless of the average value of blood pressure during the day (7). Blood pressure regulators (adrenaline, noradrenaline, cortisol, vagus tone and heart rate) are also under circadian rhythm regulation. Burgess et al. compared the autonomic nervous system function in subjects who slept through the night and those who were sleep deprived. It was observed that the rhythms of the sympathetic tone did not depend on sleep, while the vagus mediated cardiac tone showed circadian rhythmicity. However, exposure to light during sleep deprivation, which could affect sympathetic activity, was not controlled (8). Dysregulated circadian rhythm may consequently result in increase of proinflammatory cytokines. Namely, it was observed that night shift workers who stay awake up to 24 hours have increase of some markers of inflammation such as IL-1 β , IL-6, TNF- α . They were also associated with impaired glucose metabolism. – resulting in higher presence of cardiovascular events in this population (9).

Melatonin-do we have a new antihypertensive agent?

Melatonin, in addition to being a sleep hormone, has also been shown to be a potential antihypertensive agent. In a double-blind, placebo-controlled study conducted on 14 healthy men, administration of 1 mg of melatonin reduced systolic, diastolic and mean arterial pressure, as well as catecholamine levels (10). In another, also double-blind, placebo-controlled study, three-week oral administration of this hormone (2.5 mg daily) one hour before bedtime in patients with previously untreated essential hypertension significantly reduced both systolic and diastolic blood pressure (11). In these studies, the difference in blood pressure when using melatonin versus placebo was about 5 to 10 mmHg, and it is known that this reduction in diastolic pressure is associated with a 20% reduction in cardiovascular mortality (12). It has been shown that melatonin achieves hypotensive effects through central and peripheral mechanisms, as well as that receptors are located in the central and peripheral nervous system, endocrine, reproductive, gastrointestinal, immune and cardiovascular systems, including coronary arteries, cerebral arteries, aorta and systemic circulation arteries, as well as the myocardial ventricles (13,14). In the body, melatonin acts through three types of receptors, of which MT1 and MT2 play a key role in cardiovascular regulation (15).

It is assumed that hypotensive effects of melatonin are achieved through endothelium mediated vasodilatory effect, antioxidant effect and sympathetic-vagal autonomic regulation. It is a strong stimulator of nitric oxide synthase and has a vasodilatory effect through the MT2 receptor. Also, it reduces the concentration of circulating noradrenaline, and increases the cardiac tone of the vagus. Melatonin has been shown to be a safe and effective additive antihypertensive agent. The melatonin receptors are widely distributed in the body and further research is needed to elucidate all precise molecular mechanisms of its antihypertensive effects (16,17).

Disruption of the circadian rhythm and the occurrence of cardiovascular events

The modern way of life with numerous stressful situations, insufficient sleep, late, irregular and excessive meals, exposure to artificial lighting, especially the blue spectrum of light emitted by mobile phones and computers, transoceanic flights with time zones changes, shift work, especially night work, introduce a new field in science and medicine (18). Social jet lag is defined as a chronic misalignment of the circadian system that alters social behavior. Approximately 70% of the population suffers from social jet lag (for example, their biological and social clocks differ by more than one hour). Research has shown that greater social jet lag carries a greater risk of developing metabolic disorders, such as diabetes mellitus and obesity, as well as mental disorders. The association between social jet lag and cardiovascular diseases has not been fully investigated, but it is known that heart rate and cortisol levels are increased in healthy subjects, while a higher value of glycosylated hemoglobin was noticed in patients with type II diabetes mellitus (19). Epidemiological studies have shown the increased incidence of cardiovascular events in the morning hours between 6 a.m. and noon because of disrupted circadian rhythms (20). The reasons for the morning occurrence of acute myocardial infarction, stroke, malignant arrhythmias, as well as sudden cardiac death are found in increased sympathetic tone, vasoconstriction, an increase in peripheral arterial resistance and a consequent raise in blood pressure and heart rate, as well as increased platelet aggregation in the early morning hours. In contrast, a decrease in thrombolytic activity and parasympathetic tone as well as endothelial dysfunction is noted (21-24).

Endothelial dysfunction

The endothelium plays a very important role in the regulation of precise homeostatic mechanisms in the cardiovascular system (for example, vascular tone, platelet aggregation, fibrinolysis) (25). Impaired endothelial function results in vasoconstriction and intravascular thrombosis. This is particularly prominent event in the early

morning hours when increased sympathetic activity and sensitivity of α -receptors are present (26). The highest platelet aggregation and the highest concentration of plasminogen activation inhibitor-1 (PAI-1), the main regulator of endogenous fibrinolysis, is in the morning. An elevated PAI-1 is associated with the risk for atherosclerosis and thrombosis (27). Similarly, there is a morning peak of glycoprotein IIb/IIIa receptors activation on the platelet surface, which are involved in the final common pathway of platelet aggregation (28). The clinical significance is that platelet aggregation stimulated by thrombin, ADP, collagen, serotonin and other mediators can be blocked with glycoprotein IIb/IIIa receptor inhibitor drugs.

The role of the circadian system and melatonin in acute myocardial infarction

Studies have shown that the onset of acute myocardial infarction is associated with reduced nocturnal melatonin secretion. Decreased secretion of melatonin during the night in patients with ST-elevation myocardial infarction (STEMI) correlated with an increase in C-reactive protein, a marker of the acute phase of the inflammatory process, and was a predictor of adverse events in the following 6 months (29). In patients with coronary disease or heart failure, a decrease in nocturnal secretion of melatonin or its metabolite 6-sulfatoxymelatonin (aMT6S), excreted via urine, was observed (30,31). On the isolated rat hearts, melatonin had a protective effect on myocardial ischemia-reperfusion damage and reduced necrosis and the size of the infarct zone (32). For the first time in the human population a study by Reiter and al. from 2011 showed that the size of the infarct zone and the ejection fraction of the left ventricle in patients with myocardial infarction with ST segment elevation have a circadian dependence. In their study, the greatest myocardial damage was recorded one hour after the onset of ischemia and five hours after reperfusion (33). Research by Mahmoud et al showed that the highest incidence of stent thrombosis was recorded in the early morning hours (34). It is interesting that clock shifting for one hour during summer time is light saving time is associated with transient increase in the incidence of acute myocardial infarction in the spring as the consequence of sleep deprivation and poorer sleep quality, while for the winter time data are controversial (35). In a prospective randomized study, Domínguez-Rodríguez et al. showed that intravenous administration of melatonin in patients with STEMI treated by primary percutaneous coronary intervention correlated with a reduction in mortality and rehospitalization due to heart failure in a two-year follow-up period. The effect of melatonin was reflected in the attenuation of matrix metalloproteinase-9, a key enzyme involved in myocardial remodeling due to acute myocardial infarction (36). A study published in Lancet in 2018 showed the importance of chronobiology in cardiac surgery. Perioperative myocardial injury due to aortic valve replacement was significantly lower in patients operated in the afternoon, compared to surgeries performed in the morning.

This could be due to circadian expression of the myocardial gene with the nuclear receptor Rev-Erba, which is the highest in the morning. Cardioprotection would be made possible by the eventual pharmacological application of Rev-Erba receptor antagonists (37).

Circadian distribution of cardiovascular events in patients with breathing disorders during sleep

Patients with heart failure with reduced ejection fraction (HFrEF) often have breathing disorders during sleep as a comorbidity, such as obstructive (OSA) and central sleep apnea (CSA). Disruption of the circadian rhythm is associated with disturbed sleep due to various breathing disorders, with the consequent imbalance of autonomic, hemodynamic, humoral and vascular regulation (38). It has been observed that sudden cardiac death occurs most often during sleep, in the period from midnight to six in the morning, in contrast to general population, where the incidence is the highest in the morning, after waking up (39). This is supported by the findings of Bitter et al, who showed that in patients with OSA and HFrEF activations of the implantable cardioverter defibrillator (ICD) due to ventricular tachycardia were registered most often during the night, from 00:00 am to 05:59 am, while in patients with CSA activations occur at any time of the day or night (40). Obstructive apnea is known to occur only during sleep, while Cheyne Stokes breathing, as a central respiratory disorder, can also occur during wakefulness (41). Episodes of apnea lead to hypoxemia, hypercapnia, increased sympathetic tone, increased intrathoracic pressure and cardiac arrhythmias (38). Therefore, a careful, multidisciplinary evaluation of these vulnerable patient groups is needed.

Conclusion

Although numerous studies have shown the impact of circadian rhythm changes on the cardiovascular system, it should be emphasized that most of them were conducted on the healthy subjects. Therefore, potential studies of circadian rhythm should be performed on people suffering from cardiovascular diseases. It is known that cardiovascular drugs are used at the certain time of the day, in order to achieve maximum effects and while minimizing side effects. Future studies would provide detailed clarification of the circadian system role in cardiovascular complications in humans, as well as a better understanding and development of chronotherapy, with possible implementation not only in medical treatment, which can be implemented not only in medical treatment, but also in elective cardiovascular interventions, with adaptation to the biological rhythm of the individual patient.

There is no conflict of interest.

Literature

1. Stanojlović O. et al. Cirkadijalni sistem: Mreža neurona sa suprahijazmatičnim jedrom na vrhu hijerarhijske organizacije. *MedPodml* 2018; 69(3): 3–13.
2. Young MW. Time Travels: A 40-Year Journey from *Drosophila*'s Clock Mutants to Human Circadian Disorders (Nobel Lecture). *Angew Chem Int Ed Engl*. 2018 Sep 3.
3. Silver R and Rainbow M. The Suprachiasmatic Nucleus and the Circadian Timekeeping System of the Body. *Neuroscience in the 21st Century*, D.W. Pfaff (ed.), 2015; 1-49.
4. Pandi-Perumal SR, Srinivasan V, Maestroni GJM, Cardinali DP, Poeggeler B, Hardeland R. Melatonin: Nature's most versatile biological signal? *The FEBS journal* 2006; 273: 2813–38.
5. Thosar SS, Butler MP, Shea SA. Role of the circadian system in cardiovascular disease. *J Clin Invest*. 2018; Jun 1; 128(6): 2157–2167.
6. Cortés-Ríos J, Rodríguez-Fernandez M. Circadian Rhythm of Blood Pressure of Dipper and Non-dipper Patients With Essential Hypertension: A Mathematical Modeling Approach. *Front Physiol*. 2021 Jan 18; 11: 536146.
7. Dolan E. et al. Superiority of ambulatory over clinic blood pressure measurement in predicting mortality: the Dublin outcome study. *Hypertension*. 2005; 46(1): 156–161.
8. Burgess HJ, Trinder J, Kim Y, Luke D. Sleep and circadian influences on cardiac autonomic nervous system activity. *Am J Physiol*. 1997; 273(4 Pt 2): H1761–H1768.
9. Morris CJ, Purvis TE, Hu K, Scheer FA. Circadian misalignment increases cardiovascular disease risk factors in humans. *Proc Natl Acad Sci U S A*. 2016; 113(10): E1402–E1411.
10. Arangino, S, Cagnacci, A, Angiolucci, M, Vacca, AMB, Longu, G, Volpe, A & Melis, GB. Effects of melatonin on vascular reactivity, catecholamine levels, and blood pressure in healthy men. *Am J Cardiol* 1999; 83, 1417.
11. Scheer, FA, Van Montfrans, GA, Van Someren, EJ, Mairuhu, G & Buijs, RM. Daily night-time melatonin reduces blood pressure in male patients with essential hypertension. *Hypertension* 2004; 43: 192–197.
12. Rich-Edwards JW, Manson JE, Hennekens CH, Buring JE. The primary prevention of coronary heart disease in women. *N Engl J Med* 1995; 332: 1758–1766.
13. Ekmekcioglu C, Thalhammer T, Humpeler S, Mehrabi MR, Glogar HD, Holzenbein T, et al. The melatonin receptor subtype MT2 is present in the human cardiovascular system. *J Pineal Res*. 2003; 35(1): 40–4.
14. Ekmekcioglu C, Haslmayer P, Philipp C, Mehrabi MR, Glogar HD, Grimm M, et al. 24h variation in the expression of the MT1 melatonin receptor subtype in coronary arteries derived from patients with coronary heart disease. *Chronobiol Int*. 2001; 18(6): 973–85.
15. Slominski RM, Reiter RJ, Schlabritz-Loutsevitch N, Ostrom RS, Slominski AT. Melatonin membrane receptors in peripheral tissues: Distribution and functions. *Mol Cell Endocrinol*. 2012; 351: 152–66.
16. Baker J, Kimpinski K. Role of melatonin in blood pressure regulation: An adjunct anti-hypertensive agent. *Clin Exp Pharmacol Physiol*. 2018 Aug; 45(8): 755–766.
17. Saini C, Brown SA, Dibner C. Human peripheral clocks: applications for studying circadian phenotypes in physiology and pathophysiology. *Front Neurol*. 2015; 6: 95.2

18. Rutters F. et al. Is social jetlag associated with an adverse endocrine, behavioral, and cardiovascular risk profile? *J Biol Rhythms*. 2014; 29(5): 377–383.
19. Reutrakul S, Hood MM, Crowley SJ, et al. Chronotype is independently associated with glycemic control in type 2 diabetes. *Diabetes Care*. 2013; 36: 2523–2529.
20. Thosar SS, Shea SA. Circadian control of human cardiovascular function. *Curr Opin Pharmacol*. 2021 Apr; 57: 89–97.
21. Virag JA, Lust RM. Circadian influences on myocardial infarction. *Front Physiol*. 2014 Oct 30; 5: 422.
22. Muller JE, et al. Circadian variation in the frequency of onset of acute myocardial infarction. *N Engl J Med*. 1985; 313(21): 1315–1322.
23. Goldberg RJ, et al. Time of onset of symptoms of acute myocardial infarction. *Am J Cardiol*. 1990; 66(2): 140–144.
24. Elliott WJ. Circadian variation in the timing of stroke onset: a meta-analysis. *Stroke*. 1998; 29(5): 992–996.
25. Otto ME, Svatikova A, Barretto RB, Santos S, Hoffmann M, Khandheria B, Somers V. Early morning attenuation of endothelial function in healthy humans. *Circulation*. 2004 Jun 1; 109(21): 2507–10.
26. Panza JA, Epstein SE, Quyyumi AA. Circadian variation in vascular tone and its relation to alpha-sympathetic vasoconstrictor activity. *Engl J Med*. 1991 Oct 3; 325(14): 986–90.
27. Andreotti F, et al. Major circadian fluctuations in fibrinolytic factors and possible relevance to time of onset of myocardial infarction, sudden cardiac death and stroke. *Am J Cardiol*. 1988; 62(9): 635–637.
28. Scheer FA, et al. The human endogenous circadian system causes greatest platelet activation during the biological morning independent of behaviors. *PLoS One*. 2011; 6(9): e24549.
29. Dominguez-Rodriguez A, Abreu-Gonzalez P, Garcia MJ, Sánchez J, Marrero F, de Armas-Trujillo D. Decreased nocturnal melatonin levels during acute myocardial infarction. *J Pineal Res* 2002; 33: 248–252.
30. Girotti, L, Lago, M, Ianovsky, O, Carbajales, J, Elizari, MV, Brusco, LI & Cardinali, DP. Low urinary 6-sulphatoxymelatonin levels in patients with coronary artery disease. *J Pineal Res* 2002; 29: 138–142.
31. Girotti, L, Lago, M, Ianovsky, O, Elizari, MV, Dini, A, Lloret, SP, Albornoz, LE & Cardinali, DP. Low urinary 6-sulphatoxymelatonin levels in patients with severe congestive heart failure. *Endocrine* 2003; 22: 245–248.
32. Petrosillo G, Colantuono G, Moro N, Ruggiero FM, Tiravanti E, Di Venosa N, et al. Melatonin protects against heart ischemia-reperfusion injury by inhibiting mitochondrial permeability transition pore opening. *Am. J. Physiol. Heart Circ. Physiol*. 2009; 297: H1487–H1493. 10.1152/ajpheart.00163.2009
33. Ronald Reiter, Cory Swingen, Luke Moore, Timothy D Henry, and Jay H. Traverse. Circadian Dependence of Infarct Size and Left-Ventricular Function Following ST-Elevation Myocardial Infarction. *Circ Res*. 2012 Jan 6; 110(1): 105–110.
34. Mahmoud KD, Lennon RJ, Ting HH, Rihal CS, Holmes DR. Circadian variation in coronary stent thrombosis. *J Am Coll Cardiol Interv*. 2011; 4: 183–90.

35. Jiddou MR, Pica M, Boura J, Qu L, Franklin BA. Incidence of myocardial infarction with shifts to and from daylight savings time. *Am J Cardiol.* 2013; 111(5): 631–635.
36. Domínguez-Rodríguez A, Hernández-Vaquero D, Abreu-González P, Báez-Ferrer N, Díaz R, Avanzas P, Simko F, Domínguez-González V, Sharma R, Reiter RJ. Early Treatment of Acute Myocardial Infarction with Melatonin: Effects on MMP-9 and Adverse Cardiac Events. *J Clin Med.* 2022 Mar 30; 11(7): 1909.
37. Montaigne D, Marechal X, Modine T, Coisne A, Mouton S, Fayad G, et al. Daytime variation of perioperative myocardial injury in cardiac surgery and its prevention by Rev-Erb α antagonism: a single-centre propensity-matched cohort study and a randomised study. *Lancet.* 2018 Jan 6; 391(10115): 59–69.
38. Omran H, Bitter T, Horstkotte D, Oldenburg O, Fox H. Characteristics and circadian distribution of cardiac arrhythmias in patients with heart failure and sleep-disordered breathing. *Clin Res Cardiol.* 2018 Oct; 107(10): 965–974.
39. Gami AS, Howard DE, Olson EJ, Somers VK. Day-night pattern of sudden death in obstructive sleep apnea. *N Engl J Med.* 2005 Mar 24; 352(12): 1206–14.
40. Bitter T, Fox H, Dimitriadis Z, Niedermeyer J, Prib N, Prinz C, Horstkotte D, Oldenburg O. Circadian variation of defibrillator shocks in patients with chronic heart failure: the impact of Cheyne-Stokes respiration and obstructive sleep apnea. *Int J Cardiol* 2014; 176: 1033–1035.
41. Grimm W, Kesper K, Cassel W, Timmesfeld N, Hildebrandt O, Koehler U. Cheyne-stokes respiration during wakefulness in patients with chronic heart failure. *Sleep Breath* 2017; 21: 419–426.